



Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

Atto Thio12-N

Category	Fluorescent Dyes
Modification Code	AttoThio12-N
Reference Catalog Number	26-6968
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	699



[Click here for a list of fluorophores.](#)

This modification is a post synthesis conjugation to a primary amino group thus an additional modification with an amino group is required. A C3, C6 or C12 amino group can be placed at the 5' or for the 3' end a C3 or C7 amino and for internal positions an amino modified base is used, e.g Amino dT C6.

Yield of Post Synthesis NHS, Maleimide & Click Ligand Conjugation* Oligo Scale of Synthesis Yield, nmols 50 nmol 2 nmol 200 nmol 5 nmol 1 umol 16 nmol 2 umol 30 nmol 5 umol 75 nmol 10 umol 150 nmol 15 umol 225 nmol * The yield will be lower for oligos longer than 50mer. Click here for yield table of long oligos. * Click here for RNA Oligos scale of synthesis and yield. **NHS Ligand conjugation** requires a primary amino group. Gene Link offers a wide selection of amino modifications for 5', 3' and internal sites. Click here for a list of conjugation chemistry modifications. **Maleimide Ligand conjugation** requires a thiol group. Gene Link offers a wide selection of thiol modifications for 5', 3' and internal sites.

Click here for a list of conjugation chemistry modifications. **Click Chemistry Ligand conjugation** requires a corresponding Click modification; examples Alkyne:Azide, Azide:DBCO, BCN:Azide, BCN: TCO:Tetrazine. Gene Link offers a wide selection of click modifications for 5', 3' and internal sites. Click here for a list of click chemistry modifications.

ATTO Thio12 is a new label closely related to the well-known rhodamines. The characteristic feature of the label is high yield of triplet formation. The dye is moderately hydrophilic. Conventional and popular dyes that are derivatives of fluorescein (FAM, HEX and TET) and Cyanine dye derivatives (Cy3, Cy5, Cy5.5, Cy7 etc) are commonly used for fluorescently labeling oligos for use as molecular probes for real time PCR, FISH analysis and fragment analysis. For most purposes these provide a good range in wavelength and other optical properties and are available as amidites for direct coupling to oligos using automated chemistry. Other fluorescent dyes are available as N-hydroxysuccinimide (NHS) for conjugation using a primary amine group linked to the oligos. A new series of Atto dyes are now available that are designed for high sensitivity applications, including single-molecule detection.

ATTO Dyes are a series of fluorescent labels and dyes manufactured by ATTO-TEC GmbH in Siegen, Germany. The ATTO Dye series covers a spectral range from 390 nm in the UV to 740 nm in the near infrared allowing excitation with most commonly used light sources. The dyes typically are derivatives of coumarins, rhodamines, carbopyronins and oxazines. Compared with other labels especially for the red region of the spectrum, ATTO-labels show excellent photostability and brightness. Atto labels have rigid structures that do not show any cis-trans isomerization. Thus these labels display exceptional intensity with minimal spectral shift on conjugation. The molecules of most common dyes, e.g. cyanines, have a more or less flexible structure. Hence their solutions contain a mixture of several isomers with varying properties. Since the equilibrium between the isomers depends on temperature and other environmental factors, absorption and fluorescence of such dyes are ill-defined. ATTO-dyes have a molecular structure that ensures high rigidity of the chromophore. They do not form equilibria with various isomers, their optical properties are nearly independent of solvent and temperature. ATTO 647N fluoresces twice as strong as Cy5 in aqueous solution. In addition many common fluorescent labels especially cyanine dyes like Cy5 deteriorate even without any irradiation (in the dark), in particular when exposed to small concentrations of ozone present in the laboratory atmosphere. Under identical conditions of ozone exposure the new dyes ATTO 633, ATTO 647N and ATTO 655 last up to 100 times longer than cyanines like Cy5 and Alexa Fluor 647. This is very important in microarray applications, where the dye molecules are located at the surface and thus are in direct contact with the atmosphere.