



Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

Fam-TEG Azide

Category	Click Chemistry
Modification Code	Fam-TEG-N3
Reference Catalog Number	26-6722
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	576.55



6-FAM-TEG Azide
[26-6722-XX]

This modification is a post synthesis conjugation to an alkyne or DBCO modification at the appropriate site for click conjugation.

6-FAM (6-carboxyfluorescein)-TEG Azide is a 6-FAM fluorescent dye attached to a 15-atom mixed polarity triethylene glycol spacer with an azide group at the end of the spacer. 6-FAM is the most commonly used fluorescent dye for labeling oligonucleotides, and is reactive and water-soluble, with an absorbance maximum of 492 nm and an emission maximum of 517 nm. The presence of the azide allows the user to use "Click Chemistry" (a [3+2] cycloaddition reaction between alkynes and azides, using copper (I) iodide as a catalyst) to conjugate the 6-FAM-TEG Azide to a terminal alkyne-modified oligo with extremely high regioselectivity and efficiency (1,2). Preparation of the alkyne-modified oligo can be achieved using the 5'-Hexynyl modifier (see its respective tech sheet for details). The spacer acts to minimize steric hindrance between the biotin moiety and the oligo. **References**

- Huisgen, R. *Angew. Chem. Int. Ed.* (1963), **2**: 565-568.
- Rostovtsev, V.V., Green, L.G., Fokin, V.V., Sharpless, K.B. A Stepwise Huisgen Cycloaddition Process: Copper(I)-Catalyzed Regioselective Ligation of Azides and Terminal Alkynes. *Angew. Chem. Int. Ed.* (2002), **41**: 2596-2599.