



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Coumarin Azide

Category	Click Chemistry
Modification Code	Cou-N3
Reference Catalog Number	26-6726
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	203.15



**Coumarin Azide**  
**[26-6726-XX]**

This modification is a post synthesis conjugation to an alkyne or DBCO modification at the appropriate site for click conjugation.

Coumarin (7-Hydroxycoumarin)-Azide is a fluorescent dye containing a terminal azide group. Coumarin is also known as umbelliferone. Coumarin is highly fluorescent and pH-sensitive, with an absorbance maximum of 358 nm and an emission maximum of 480 nm; thus it emits in the blue region of the visible spectrum. The presence of the azide allows the user to use "Click Chemistry" (a [3+2] cycloaddition reaction between alkynes and azides, using copper (I) iodide as a catalyst) to conjugate the Coumarin-Azide to a terminal alkyne-modified oligo with extremely high regioselectivity and efficiency (1,2). Preparation of the alkyne-modified oligo can be achieved using the 5'-Hexynyl modifier (see its respective tech sheet for details). Because coumarin is effectively quenched if its hydroxyl group is either alkylated or phosphorylated, it is useful in high-throughput screening for enzyme lipases and phosphatases. **References**

- Huisgen, R. *Angew. Chem. Int. Ed.* (1963), **2**: 565-568.
- Rostovtsev, V.V., Green, L.G., Fokin, V.V., Sharpless, K.B. A Stepwise Huisgen Cycloaddition Process: Copper(I)-Catalyzed Regioselective Ligation of Azides and Terminal Alkynes. *Angew. Chem. Int. Ed.* (2002), **41**: 2596-2599.