



## Product Specifications

Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates  
Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

## FISH Probes Introduction

Probes for fluorescent in situ hybridization (FISH) are derived from repetitive chromosomal regions to paint a chromosome or are directed towards unique gene(s) for identification. These probes are usually developed from clones (BAC and cosmids) and more recently synthesized chemically as oligonucleotides appropriately labeled with fluorescent dyes. The oligo probes can be synthesized as tiles or longer sequences up to 250mer in length by Gene Link. Gene Link presents various design options for synthesizing 'Smart' FISH probes. Smart FISH probes can be synthesized in a predetermined way to exhibit the features that is desired; for instance to increase nuclease resistance we can substitute standard bases with 2' Fluoro bases. Standard DNA probes are like natural molecules that are prone to degradation under normal conditions, ubiquitous nucleases as well as chemical instability lead to fast degradation with a finite half life. The premise of this product guide is to discuss various modifications that are offered by Gene Link that may be used for design and synthesis of Smart FISH Probes. There are four common features that are desirable and in particular can be improved by using a combination of available nucleic acid modifications that modify the phosphodiester linkages, nucleic acid bases, the sugar moieties and addition of various other functional groups.

# FISH Probes Design Protocols

## Design Guidelines for FISH Probes

1. Design multiple 24 to 30mer probes. Avoid stretches of more than 3 G or C bases.
2. To impart exonuclease resistance substitute 3-4 bases at the 5' and 3' end with 2'F bases. The 2' F bases imparts resistance to exonuclease degradation and increases duplex stability by 4-6 degrees.
3. Several internal bases can be substituted with 5me dC and 2 Amino dA to further increase duplex stability.
4. Affinity ligands such as Digoxigenin or Biotin or fluorescent dye e.g Cy3, Cy5 or any other can be labeled at the 3' and 5' end. Multiple internal sites can also be labeled with affinity ligands or fluorescent dyes to increase sensitivity.
5. Multiple dye sites should be spaced apart by 10 or more bases.
6. The above guidelines are for all initial FISH probe design. Design rules may have to be established empirically for very specific or novel assay settings, but following the above recommendations will provide a good start.

## FISH Probes **Applications**

Probes for fluorescent in situ hybridization (FISH) are derived from repetitive chromosomal regions to paint a chromosome or are directed towards unique gene(s) for identification. These probes are usually developed from clones (BAC and cosmids) and more recently synthesized chemically as oligonucleotides appropriately labeled with fluorescent dyes. The oligo probes can be synthesized as tiles or longer sequences up to 250mer in length by Gene Link.

## References

1. Milligan, J.F., Matteucci, M.D. and Martin, J.C. (1993) Current concepts in antisense drug design. *J. Medicinal Chem.* 36:1923-1937.
2. Helene, C., Toulme, J. (1990) Specific regulation of gene expression by antisense, sense and antigene nucleic acids. *Biochim. Biophys. Acta.* 1049: 99-125.
3. Weintraub, H. M. (1990) Antisense RNA and DNA. *Sci. Amer.* 262:40-46.
4. Iyer, R.P., Egan, W., Regan, J.B and Beaucage, S.L. (1990) *J. Am. Chem. Soc.* 112; 1253-1254.
5. Wagner, R.W., Matteucci, M.D., Lewis, J.G., Gutierrez, A.J., Moulds, C. and Froehler, B.C. (1993) Antisense gene inhibition by oligonucleotides containing C-5 propyne pyrimidines. *Science* 260:1510-1513.
6. Hertoghs, K.M.L., Ellis, J.H. and Catchpole, I.R (2003) Use of locked nucleic acid oligonucleotides to add functionality to plasmid DNA. *Nucl. Acids Res.* 31 (20): 5817-5830.
7. Cotton, M., Oberhauser, B., Burnar, H. et al. (1991) 2'O methyl and 2'O ethyl oligoribonucleotides as inhibitors of the in vitro U7 snRNPdependent messenger-RNA processing event. *NAR* 19:2629.
8. Singh, S.K., Nielsen, P., Koshkin, A.A. and J. Wengel, *Chem. Comm.*, 1998, (4), 455-456.
9. A.A. Koshkin, A.A., Singh, S.K., Nielsen, P., Rajwanshi, V.K., Kumar, R., Meldgaard, M., Olsen, C.E and Wengel, J. (1998) *Tetrahedron* 54:3607-3630.
10. Kværnø, L. and Wengel, J. (1999) *Chem. Comm.*, 7:657-658.
11. Petersen, M and Wengel, J. (2003) *Trends in Biotechnology* 21(2): 74-81.
12. P.A. Giannaris, P.A. and Damha, M.J (1993) *Nucleic Acids Research*, 21:4742-4749.
13. Bhan, P., Bhan, A., Hong, M.K., Hartwell, J.G., Saunders, J.M and Hoke, G.D (1997) *Nucleic Acids Res*, 25:3310-3317.
14. Coleman, R.S. and Kesicki, E.S (1994) *J. Amer. Chem. Soc.*, 116:11636-11642. 15. John SantaLucia, Jr. (1998) *Proc. Natl. Acad. Sci.* 95 1460-1465.

## Modificaton Code List

Modification	Code	Catalog Number
2'-Fluoro deoxyadenosine (2'-F-A)	[fA]	26-6692
2'-Fluoro deoxycytosine (2'-F-C)	[fC]	26-6463
2'-Fluoro deoxyguanosine (2'-F-G)	[fG]	26-6693
2'-Fluoro deoxyuridine (2'-F-U)	[fU]	26-6462
2'-O Me-5-Me-C	[m5mC]	27-6508
2'-O methyl adenosine A	[mA]	27-6410A
2'-O methyl cytosine C	[mC]	27-6410C
2'-O methyl guanosine G	[mG]	27-6410G
2'-O methyl uridine U	[mU]	27-6410U
2-Amino dA	[2Am-dA]	26-6525
5-methyl deoxycytosine [5mdC]	[5mdC]	26-6413
Amino C6 U (RNA)	[AmC6U]	27-6422
Amino deoxyadenosine dA C6	[Am-dA-C6]	26-6666
Amino dC N4 PEG3 amino linker	[Am-dC-C6]	26-6670
Amino deoxyguanosine dG C6	[Am-dG-C6]	26-6669
Amino deoxythymidine dT C6	[Am-dT-C6]	26-6438
Cy3-5' (Cyanine 3)	[Cy3-5]	26-6437
Cy3.5 (Cyanine 3.5-5')	[Cy3.5-5]	26-6461
Cy5 (5'-Cyanine 5)	[Cy5-5]	26-6436
Cy5.5 (Cyanine 5.5)	[Cy5.5-5]	26-6460

Cy7 (Cyanine 7)	[Cy7-N]	26-6474
Fluorescein dT (Fam dT)	[Fam-dT]	26-6422
Fluorescein-5' (FITC)	[FI-5]	26-6426
Hex-5' (Hexachloro-Fluorescein)	[Hex-5]	26-6432
Tet-5' (Tetrachloro-Fluorescein)	[Tet-5]	26-6433



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 2'-F-A

Category	Nuclease Resistance
Modification Code	fA
Reference Catalog Number	26-6692
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	331.2



Antisense Oligos (ODN) & siRNA Oligo Modifications

**Click here for more information on antisense modifications, design & applications.**

2'-Fluoro-deoxyadenosine (2'-F-A) is classified as a 2'-Fluoro RNA monomer. The substitution of a highly electronegative fluorine atom for the 2'-OH group of the ribose results in the ring of a 2'-F-ribonucleoside having a C3'-endo (i.e., RNA-type) conformation. Consequently, an RNA oligonucleotide containing a 2'-F RNA monomer adopts the more thermodynamically stable A-form helix on hybridization to a target (1).

2'-F RNA modifications have been shown to be particularly useful in the following oligo-based applications:

(a) **Anti-sense oligos & Nuclease Resistance:** When 2'-F RNA is incorporated into an anti-sense DNA oligo (resulting in a 2'-F RNA/DNA chimeric), the  $T_m$  of its duplex with RNA **increases** relative to that formed by an unmodified anti-sense DNA oligo **by 1.8°C per 2'-F RNA residue added** (1). The DNA/RNA duplex formed between a chimeric 2'-F-RNA/DNA anti-sense oligo and its RNA target also is a substrate for RNase H. With respect to nuclease resistance, while 2'-F RNA nucleotides do provide some nuclease resistance when incorporated into DNA, phosphorothiolation of the 2'-F RNA phosphodiester linkages is recommended, because it strongly enhances such resistance. This effect becomes particularly important if the 2'-F RNA nucleotide is to be incorporated at one or more of the first three positions of the 5'- or 3'-ends of the oligo.

Modifications Increasing Duplex Stability and Nuclease Resistance

Modification

Duplex Stability [ $T_m$  Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.

7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased Click here for complete list of duplex stability modifications

(b) **Aptamers:** The addition of 2'-F pyrimidines (C, U) into RNA aptamers has been shown to provide them with both considerably increased nuclease resistance and equal or higher binding affinity for their ligands, compared to the corresponding unmodified aptamers (2,3). The incorporation of 2'-F RNA nucleotides into RNA aptamers as an optimization strategy is becoming an increasingly common practice.

(c) **siRNA & Nuclease Resistance:** siRNA synthesized with 2'-F pyrimidines have been shown to have greatly increased stability in plasma compared to 2'-OH siRNA (4,5). In one study, levels of inhibition for 2'-F siRNA and 2'-OH siRNA, in cell culture and *in vivo* using BALB/c mice transfected with pGL3 luciferase, were similar over time (4). In another study, siRNA fully substituted with both 2'-F RNA and 2'-O-Methyl RNA nucleotides displayed not only enhanced stability in plasma, but also >500-fold increase in capability to down-regulate gene expression compared with the corresponding unmodified siRNA (5).

(d) **LNA Alternative:** The increased thermal stability and nuclease resistance provided by 2'-F RNA residues make them attractive as lower-cost substitutes in any application involving LNA-modified oligos.

#### **ASO's and siRNA Modifications.**

Click this link to view ASO's and siRNA Modifications.

**ASO's and siRNA Delivery.** The development of effective delivery systems for antisense oligonucleotides is essential for their clinical therapeutic application. The most common delivery system involves a relatively hydrophobic molecule that can cross the lipid membrane. Cholesterol TEG, alpha-Tocopherol TEG ( a natural isomer of vitamin E), stearyl and GalNAc modifications have been shown to effective for delivery of ASO's and siRNA in addition to cell penetrating peptides. Click this link to view these modifications.

#### **REFERENCES:**

1. Kawasaki, A.M., et al., Uniformly modified 2'-deoxy-2'-fluoro phosphorothioate oligonucleotides as nuclease resistant antisense compounds with high affinity and specificity for RNA targets, *Journal of Medicinal Chemistry* (1993), **36**: 831-841.
2. Khati, M.; Schuman, M.; Ibrahim, J.; Sattentau, Q.; Gordon, S.; James, W. Neutralization of infectivity of diverse R5 clinical isolates of human immunodeficiency virus type 1 by gp120-binding 2'-F-RNA aptamers. *J Virol*, (2003). **77**: 2692-8.
3. Goring, H.U.; Adler, Annette; Forster, Nicole; Homann, Matthias. Post-SELEX Chemical Optimization of a Trypanosome-Specific RNA Aptamer. *Combinatorial Chemistry & High Throughput Screening* (2008), **11**: 16-23.
4. Layzer, J.M.; McCaffrey, A.P.; Tanner, A.K.; Huang, Z.; Kay, M.A.; Sullenger, B.A. In vivo activity of nuclease-resistant siRNAs. *RNA* (2004), **10**, 766-71.
5. Allerson, C.R.; et al. Fully 2'-modified oligonucleotide duplexes with improved in vitro potency and stability compared to unmodified small interfering RNA. *Journal of Medicinal Chemistry* (2005), **48**: 901-904.





## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 2'-F-C

Category	Nuclease Resistance
Modification Code	fC
Reference Catalog Number	26-6463
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	307.17



Antisense Oligos (ODN) & siRNA Oligo Modifications

**Click here for more information on antisense modifications, design & applications.**

2'-Fluoro-cytosine (2'-F-C) is classified as a 2'-Fluoro RNA monomer. The substitution of a highly electronegative fluorine atom for the 2'-OH group of the ribose results in the ring of a 2'-F-ribonucleoside having a C3'-endo (i.e., RNA-type) conformation. Consequently, an RNA oligonucleotide containing a 2'-F RNA monomer adopts the more thermodynamically stable A-form helix on hybridization to a target (1).

2'-F RNA modifications have been shown to be particularly useful in the following oligo-based applications:

(a) **Anti-sense oligos & Nuclease Resistance:** When 2'-F RNA is incorporated into an anti-sense DNA oligo (resulting in a 2'-F RNA/DNA chimeric), the  $T_m$  of its duplex with RNA **increases** relative to that formed by an unmodified anti-sense DNA oligo **by 1.8°C per 2'-F RNA residue added** (1). The DNA/RNA duplex formed between a chimeric 2'-F-RNA/DNA anti-sense oligo and its RNA target also is a substrate for RNase H. With respect to nuclease resistance, while 2'-F RNA nucleotides do provide some nuclease resistance when incorporated into DNA, phosphorothiolation of the 2'-F RNA phosphodiester linkages is recommended, because it strongly enhances such resistance. This effect becomes particularly important if the 2'-F RNA nucleotide is to be incorporated at one or more of the first three positions of the 5'- or 3'-ends of the oligo.

Modifications Increasing Duplex Stability and Nuclease Resistance

Modification

Duplex Stability [ $T_m$  Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.

7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased Click here for complete list of duplex stability modifications

(b) **Aptamers:** The addition of 2'-F pyrimidines (C, U) into RNA aptamers has been shown to provide them with both considerably increased nuclease resistance and equal or higher binding affinity for their ligands, compared to the corresponding unmodified aptamers (2,3). The incorporation of 2'-F RNA nucleotides into RNA aptamers as an optimization strategy is becoming an increasingly common practice.

(c) **siRNA & Nuclease Resistance:** siRNA synthesized with 2'-F pyrimidines have been shown to have greatly increased stability in plasma compared to 2'-OH siRNA (4,5). In one study, levels of inhibition for 2'-F siRNA and 2'-OH siRNA, in cell culture and *in vivo* using BALB/c mice transfected with pGL3 luciferase, were similar over time (4). In another study, siRNA fully substituted with both 2'-F RNA and 2'-O-Methyl RNA nucleotides displayed not only enhanced stability in plasma, but also >500-fold increase in capability to down-regulate gene expression compared with the corresponding unmodified siRNA (5).

(d) **LNA Alternative:** The increased thermal stability and nuclease resistance provided by 2'-F RNA residues make them attractive as lower-cost substitutes in any application involving LNA-modified oligos.

#### **ASO's and siRNA Modifications.**

Click this link to view ASO's and siRNA Modifications.

**ASO's and siRNA Delivery.** The development of effective delivery systems for antisense oligonucleotides is essential for their clinical therapeutic application. The most common delivery system involves a relatively hydrophobic molecule that can cross the lipid membrane. Cholesterol TEG, alpha-Tocopherol TEG ( a natural isomer of vitamin E), stearyl and GalNAc modifications have been shown to effective for delivery of ASO's and siRNA in addition to cell penetrating peptides. Click this link to view these modifications.

#### **REFERENCES:**

1. Kawasaki, A.M., et al., Uniformly modified 2'-deoxy-2'-fluoro phosphorothioate oligonucleotides as nuclease resistant antisense compounds with high affinity and specificity for RNA targets, *Journal of Medicinal Chemistry* (1993), **36**: 831-841.
2. Khati, M.; Schuman, M.; Ibrahim, J.; Sattentau, Q.; Gordon, S.; James, W. Neutralization of infectivity of diverse R5 clinical isolates of human immunodeficiency virus type 1 by gp120-binding 2'-F-RNA aptamers. *J Virol*, (2003). **77**: 2692-8.
3. Goring, H.U.; Adler, Annette; Forster, Nicole; Homann, Matthias. Post-SELEX Chemical Optimization of a Trypanosome-Specific RNA Aptamer. *Combinatorial Chemistry & High Throughput Screening* (2008), **11**: 16-23.
4. Layzer, J.M.; McCaffrey, A.P.; Tanner, A.K.; Huang, Z.; Kay, M.A.; Sullenger, B.A. In vivo activity of nuclease-resistant siRNAs. *RNA* (2004), **10**, 766-71.
5. Allerson, C.R.; et al. Fully 2'-modified oligonucleotide duplexes with improved in vitro potency and stability compared to unmodified small interfering RNA. *Journal of Medicinal Chemistry* (2005), **48**: 901-904.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 2'-F-G

Category	Nuclease Resistance
Modification Code	fG
Reference Catalog Number	26-6693
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	347.16



Antisense Oligos (ODN) & siRNA Oligo Modifications

**Click here for more information on antisense modifications, design & applications.**

2'-Fluoro-deoxyguanosine (2'-F-G) is classified as a 2'-Fluoro RNA monomer. The substitution of a highly electronegative fluorine atom for the 2'-OH group of the ribose results in the ring of a 2'-F-ribonucleoside having a C3'-endo (i.e., RNA-type) conformation. Consequently, an RNA oligonucleotide containing a 2'-F RNA monomer adopts the more thermodynamically stable A-form helix on hybridization to a target (1).

2'-F RNA modifications have been shown to be particularly useful in the following oligo-based applications:

(a) **Anti-sense oligos & Nuclease Resistance:** When 2'-F RNA is incorporated into an anti-sense DNA oligo (resulting in a 2'-F RNA/DNA chimeric), the  $T_m$  of its duplex with RNA **increases** relative to that formed by an unmodified anti-sense DNA oligo **by 1.8°C per 2'-F RNA residue added** (1). The DNA/RNA duplex formed between a chimeric 2'-F-RNA/DNA anti-sense oligo and its RNA target also is a substrate for RNase H. With respect to nuclease resistance, while 2'-F RNA nucleotides do provide some nuclease resistance when incorporated into DNA, phosphorothiolation of the 2'-F RNA phosphodiester linkages is recommended, because it strongly enhances such resistance. This effect becomes particularly important if the 2'-F RNA nucleotide is to be incorporated at one or more of the first three positions of the 5'- or 3'-ends of the oligo.

Modifications Increasing Duplex Stability and Nuclease Resistance

Modification

Duplex Stability [ $T_m$  Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.

7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased Click here for complete list of duplex stability modifications

(b) **Aptamers:** The addition of 2'-F pyrimidines (C, U) into RNA aptamers has been shown to provide them with both considerably increased nuclease resistance and equal or higher binding affinity for their ligands, compared to the corresponding unmodified aptamers (2,3). The incorporation of 2'-F RNA nucleotides into RNA aptamers as an optimization strategy is becoming an increasingly common practice.

(c) **siRNA & Nuclease Resistance:** siRNA synthesized with 2'-F pyrimidines have been shown to have greatly increased stability in plasma compared to 2'-OH siRNA (4,5). In one study, levels of inhibition for 2'-F siRNA and 2'-OH siRNA, in cell culture and *in vivo* using BALB/c mice transfected with pGL3 luciferase, were similar over time (4). In another study, siRNA fully substituted with both 2'-F RNA and 2'-O-Methyl RNA nucleotides displayed not only enhanced stability in plasma, but also >500-fold increase in capability to down-regulate gene expression compared with the corresponding unmodified siRNA (5).

(d) **LNA Alternative:** The increased thermal stability and nuclease resistance provided by 2'-F RNA residues make them attractive as lower-cost substitutes in any application involving LNA-modified oligos.

#### **ASO's and siRNA Modifications.**

Click this link to view ASO's and siRNA Modifications.

**ASO's and siRNA Delivery.** The development of effective delivery systems for antisense oligonucleotides is essential for their clinical therapeutic application. The most common delivery system involves a relatively hydrophobic molecule that can cross the lipid membrane. Cholesterol TEG, alpha-Tocopherol TEG ( a natural isomer of vitamin E), stearyl and GalNAc modifications have been shown to effective for delivery of ASO's and siRNA in addition to cell penetrating peptides. Click this link to view these modifications.

#### **REFERENCES:**

1. Kawasaki, A.M., et al., Uniformly modified 2'-deoxy-2'-fluoro phosphorothioate oligonucleotides as nuclease resistant antisense compounds with high affinity and specificity for RNA targets, *Journal of Medicinal Chemistry* (1993), **36**: 831-841.
2. Khati, M.; Schuman, M.; Ibrahim, J.; Sattentau, Q.; Gordon, S.; James, W. Neutralization of infectivity of diverse R5 clinical isolates of human immunodeficiency virus type 1 by gp120-binding 2'-F-RNA aptamers. *J Virol*, (2003). **77**: 2692-8.
3. Goring, H.U.; Adler, Annette; Forster, Nicole; Homann, Matthias. Post-SELEX Chemical Optimization of a Trypanosome-Specific RNA Aptamer. *Combinatorial Chemistry & High Throughput Screening* (2008), **11**: 16-23.
4. Layzer, J.M.; McCaffrey, A.P.; Tanner, A.K.; Huang, Z.; Kay, M.A.; Sullenger, B.A. In vivo activity of nuclease-resistant siRNAs. *RNA* (2004), **10**, 766-71.
5. Allerson, C.R.; et al. Fully 2'-modified oligonucleotide duplexes with improved in vitro potency and stability compared to unmodified small interfering RNA. *Journal of Medicinal Chemistry* (2005), **48**: 901-904.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 2'-F-U

Category	Nuclease Resistance
Modification Code	fU
Reference Catalog Number	26-6462
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	308.16



Antisense Oligos (ODN) & siRNA Oligo Modifications

**Click here for more information on antisense modifications, design & applications.**

2'-Fluoro-deoxyuridine (2'-F-U) is classified as a 2'-Fluoro RNA monomer. The substitution of a highly electronegative fluorine atom for the 2'-OH group of the ribose results in the ring of a 2'-F-ribonucleoside having a C3'-endo (i.e., RNA-type) conformation. Consequently, an RNA oligonucleotide containing a 2'-F RNA monomer adopts the more thermodynamically stable A-form helix on hybridization to a target (1).

2'-F RNA modifications have been shown to be particularly useful in the following oligo-based applications:

(a) **Anti-sense oligos & Nuclease Resistance:** When 2'-F RNA is incorporated into an anti-sense DNA oligo (resulting in a 2'-F RNA/DNA chimeric), the  $T_m$  of its duplex with RNA **increases** relative to that formed by an unmodified anti-sense DNA oligo **by 1.8°C per 2'-F RNA residue added** (1). The DNA/RNA duplex formed between a chimeric 2'-F-RNA/DNA anti-sense oligo and its RNA target also is a substrate for RNase H. With respect to nuclease resistance, while 2'-F RNA nucleotides do provide some nuclease resistance when incorporated into DNA, phosphorothiolation of the 2'-F RNA phosphodiester linkages is recommended, because it strongly enhances such resistance. This effect becomes particularly important if the 2'-F RNA nucleotide is to be incorporated at one or more of the first three positions of the 5'- or 3'-ends of the oligo.

Modifications Increasing Duplex Stability and Nuclease Resistance

Modification

Duplex Stability [ $T_m$  Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.

7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased [Click here for complete list of duplex stability modifications](#)

(b) **Aptamers:** The addition of 2'-F pyrimidines (C, U) into RNA aptamers has been shown to provide them with both considerably increased nuclease resistance and equal or higher binding affinity for their ligands, compared to the corresponding unmodified aptamers (2,3). The incorporation of 2'-F RNA nucleotides into RNA aptamers as an optimization strategy is becoming an increasingly common practice.

(c) **siRNA & Nuclease Resistance:** siRNA synthesized with 2'-F pyrimidines have been shown to have greatly increased stability in plasma compared to 2'-OH siRNA (4,5). In one study, levels of inhibition for 2'-F siRNA and 2'-OH siRNA, in cell culture and *in vivo* using BALB/c mice transfected with pGL3 luciferase, were similar over time (4). In another study, siRNA fully substituted with both 2'-F RNA and 2'-O-Methyl RNA nucleotides displayed not only enhanced stability in plasma, but also >500-fold increase in capability to down-regulate gene expression compared with the corresponding unmodified siRNA (5).

(d) **LNA Alternative:** The increased thermal stability and nuclease resistance provided by 2'-F RNA residues make them attractive as lower-cost substitutes in any application involving LNA-modified oligos.

#### **ASO's and siRNA Modifications.**

[Click this link to view ASO's and siRNA Modifications.](#)

**ASO's and siRNA Delivery.** The development of effective delivery systems for antisense oligonucleotides is essential for their clinical therapeutic application. The most common delivery system involves a relatively hydrophobic molecule that can cross the lipid membrane. Cholesterol TEG, alpha-Tocopherol TEG ( a natural isomer of vitamin E), stearyl and GalNAc modifications have been shown to effective for delivery of ASO's and siRNA in addition to cell penetrating peptides. [Click this link to view these modifications.](#)

#### **REFERENCES:**

1. Kawasaki, A.M., et al., Uniformly modified 2'-deoxy-2'-fluoro phosphorothioate oligonucleotides as nuclease resistant antisense compounds with high affinity and specificity for RNA targets, *Journal of Medicinal Chemistry* (1993), **36**: 831-841.
2. Khati, M.; Schuman, M.; Ibrahim, J.; Sattentau, Q.; Gordon, S.; James, W. Neutralization of infectivity of diverse R5 clinical isolates of human immunodeficiency virus type 1 by gp120-binding 2'-F-RNA aptamers. *J Virol*, (2003). **77**: 2692-8.
3. Goring, H.U.; Adler, Annette; Forster, Nicole; Homann, Matthias. Post-SELEX Chemical Optimization of a Trypanosome-Specific RNA Aptamer. *Combinatorial Chemistry & High Throughput Screening* (2008), **11**: 16-23.
4. Layzer, J.M.; McCaffrey, A.P.; Tanner, A.K.; Huang, Z.; Kay, M.A.; Sullenger, B.A. In vivo activity of nuclease-resistant siRNAs. *RNA* (2004), **10**, 766-71.
5. Allerson, C.R.; et al. Fully 2'-modified oligonucleotide duplexes with improved in vitro potency and stability compared to unmodified small interfering RNA. *Journal of Medicinal Chemistry* (2005), **48**: 901-904.





## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 2'-O Me-5-Me-C

Category	Structural Studies
Modification Code	m5mC
Reference Catalog Number	27-6508
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	333.24



Antisense Oligos (ODN) & siRNA Oligo Modifications

**Click here for more information on antisense modifications, design & applications.**

2'-OMethyl-5-methyl cytosine (2'-OMe-5-Me-C) is an RNA monomer that pairs with G, and when substituted for C in an oligonucleotide, both increases the stability of the resulting duplex relative to the comparable unmodified form, and confers nuclease resistance at that position(1). This "double-methylated"-modified cytosine thus is an excellent choice for incorporation into anti-sense oligos, where both properties are particularly desirable. Furthermore, because anti-sense oligonucleotides containing a CpG motif are known to induce pro-inflammatory responses after *in vivo* administration to animals, including human, via activation of Toll-like receptor 9 (TLR9), substitution of 2'-OMe-5-Me-dC for C in these motifs can prevent or sharply reduce these undesirable immune responses (2,3).

Modifications Increasing Duplex Stability and Nuclease Resistance

Modification

Duplex Stability [Tm Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased

asp?mod\_sp\_cat\_id=19 >Click here for complete list of duplex stability modifications **References**

1. Bundock, P.; de Both, M.T.J.; Hogers, R.C.J. 2006. Alternative nucleotides for improved targeted nucleotide exchange. Patent No. 2007073149, filed Dec 22, 2005, issued June 28, 2007.
2. Henry, S.P.; Stecker, K.; Brooks, D.; Monteith, D.; Conklin, B.; Bennett, C.F. Chemically modified oligonucleotides exhibit decreased immune stimulation in mice. *J. Pharmacol. Exp. Ther.* (2000), **292**: 468-479.
3. Yu, D.; Wang, D.; Zhu, F.-G.; Bhagat, L.; Dai, M.; Kandimalia, E.R.; Agrawal, S. Modifications Incorporated in CpG Motifs of Oligodeoxynucleotides Lead to Antagonist Activity of Toll-like Receptors 7 and 9. *J. Med. Chem.* (2009), **52**: 5108-5114.





## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 2'-O methyl A

Category	Nuclease Resistance
Modification Code	mA
Reference Catalog Number	27-6410A
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	343.24



Antisense Oligos (ODN) & siRNA Oligo Modifications

**Click here for more information on antisense modifications, design & applications.**

2'-O-Methyl bases are classified as a 2'-O-Methyl RNA monomer. 2'-O-Methyl nucleotides are most commonly used **to confer nuclease resistance** to an oligo designed for anti-sense, siRNA or aptamer-based research, diagnostic or therapeutic purposes, when specific 2'-OH is not required. Nuclease resistance can be further enhanced by phosphorothiolation of appropriate internucleotide linkages within the oligo.

The hydrogen bonding behavior of a 2'-O-Methyl RNA/RNA base pair is closer to that of an RNA/RNA base pair than a DNA/RNA base pair. Consequently, the presence of 2'-O-Methyl nucleotides **improves duplex stability**. Indeed, incorporation of a 2'-O-Methyl nucleotide into an anti-sense oligo (resulting in a 2'-O-Methyl RNA/DNA chimeric), lead to a **increase** in the T<sub>m</sub> of its duplex with RNA, relative to that formed by an unmodified anti-sense DNA oligo, **of 1.3°C per 2'-O-Methyl RNA residue added** (2). Moreover, from a synthesis standpoint, the coupling efficiency of 2'-O-Methyl phosphoramidites are higher than those of RNA monomers, resulting in higher yield of full-length oligos.

Modifications Increasing Duplex Stability and Nuclease Resistance

Modification

Duplex Stability [T<sub>m</sub> Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased

[genelink.com/oligo\\_modifications\\_reference/OMR\\_mod\\_category\\_intro.asp?mod\\_sp\\_cat\\_id=19](http://genelink.com/oligo_modifications_reference/OMR_mod_category_intro.asp?mod_sp_cat_id=19) >Click here for complete list of duplex stability modifications

### **ASO's and siRNA Modifications.**

Click this link to view ASO's and siRNA Modifications.

**ASO's and siRNA Delivery.** The development of effective delivery systems for antisense oligonucleotides is essential for their clinical therapeutic application. The most common delivery system involves a relatively hydrophobic molecule that can cross the lipid membrane. Cholesterol TEG, alpha-Tocopherol TEG ( a natural isomer of vitamin E), stearyl and GalNAc modifications have been shown to effective for delivery of ASO's and siRNA in addition to cell penetrating peptides. Click this link to view these modifications.

### **References**

1. Cotton, M.; Oberhauser, B.; Burnar, H. *et al.* 2'O methyl and 2'O ethyl oligoribonucleotides as inhibitors of the in vitro U7 snRNP-dependent messenger-RNA processing event. *Nucleic Acids Res.* (1991) , **19**:2629-2635.
2. Kawasaki, A.M. *et al.*, Uniformly modified 2'-deoxy-2'-fluoro phosphorothioate oligonucleotides as nuclease resistant antisense compounds with high affinity and specificity for RNA targets, *Journal of Medicinal Chemistry* (1993), **36**: 831-841.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 2'-O methyl C

Category	Nuclease Resistance
Modification Code	mC
Reference Catalog Number	27-6410C
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	319.21



Antisense Oligos (ODN) & siRNA Oligo Modifications

**Click here for more information on antisense modifications, design & applications.**

2'-O-Methyl bases are classified as a 2'-O-Methyl RNA monomer. 2'-O-Methyl nucleotides are most commonly used **to confer nuclease resistance** to an oligo designed for anti-sense, siRNA or aptamer-based research, diagnostic or therapeutic purposes, when specific 2'-OH is not required. Nuclease resistance can be further enhanced by phosphorothiolation of appropriate internucleotide linkages within the oligo.

The hydrogen bonding behavior of a 2'-O-Methyl RNA/RNA base pair is closer to that of an RNA/RNA base pair than a DNA/RNA base pair. Consequently, the presence of 2'-O-Methyl nucleotides **improves duplex stability**. Indeed, incorporation of a 2'-O-Methyl nucleotide into an anti-sense oligo (resulting in a 2'-O-Methyl RNA/DNA chimeric), lead to a **increase** in the  $T_m$  of its duplex with RNA, relative to that formed by an unmodified anti-sense DNA oligo, **of 1.3°C per 2'-O-Methyl RNA residue added** (2). Moreover, from a synthesis standpoint, the coupling efficiency of 2'-O-Methyl phosphoramidites are higher than those of RNA monomers, resulting in higher yield of full-length oligos.

Modifications Increasing Duplex Stability and Nuclease Resistance

Modification

Duplex Stability [ $T_m$  Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased

[genelink.com/oligo\\_modifications\\_reference/OMR\\_mod\\_category\\_intro.asp?mod\\_sp\\_cat\\_id=19](http://genelink.com/oligo_modifications_reference/OMR_mod_category_intro.asp?mod_sp_cat_id=19) >Click here for complete list of duplex stability modifications

### **ASO's and siRNA Modifications.**

Click this link to view ASO's and siRNA Modifications.

**ASO's and siRNA Delivery.** The development of effective delivery systems for antisense oligonucleotides is essential for their clinical therapeutic application. The most common delivery system involves a relatively hydrophobic molecule that can cross the lipid membrane. Cholesterol TEG, alpha-Tocopherol TEG ( a natural isomer of vitamin E), stearyl and GalNAc modifications have been shown to effective for delivery of ASO's and siRNA in addition to cell penetrating peptides. Click this link to view these modifications.

### **References**

1. Cotton, M.; Oberhauser, B.; Burnar, H. *et al.* 2'O methyl and 2'O ethyl oligoribonucleotides as inhibitors of the in vitro U7 snRNP-dependent messenger-RNA processing event. *Nucleic Acids Res.* (1991) , **19**:2629-2635.
2. Kawasaki, A.M. *et al.*, Uniformly modified 2'-deoxy-2'-fluoro phosphorothioate oligonucleotides as nuclease resistant antisense compounds with high affinity and specificity for RNA targets, *Journal of Medicinal Chemistry* (1993), **36**: 831-841.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 2'-O methyl G

Category	Nuclease Resistance
Modification Code	mG
Reference Catalog Number	27-6410G
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	359.24



Antisense Oligos (ODN) & siRNA Oligo Modifications

**Click here for more information on antisense modifications, design & applications.**

2'-O-Methyl bases are classified as a 2'-O-Methyl RNA monomer. 2'-O-Methyl nucleotides are most commonly used **to confer nuclease resistance** to an oligo designed for anti-sense, siRNA or aptamer-based research, diagnostic or therapeutic purposes, when specific 2'-OH is not required. Nuclease resistance can be further enhanced by phosphorothiolation of appropriate internucleotide linkages within the oligo.

The hydrogen bonding behavior of a 2'-O-Methyl RNA/RNA base pair is closer to that of an RNA/RNA base pair than a DNA/RNA base pair. Consequently, the presence of 2'-O-Methyl nucleotides **improves duplex stability**. Indeed, incorporation of a 2'-O-Methyl nucleotide into an anti-sense oligo (resulting in a 2'-O-Methyl RNA/DNA chimeric), lead to a **increase** in the  $T_m$  of its duplex with RNA, relative to that formed by an unmodified anti-sense DNA oligo, **of 1.3°C per 2'-O-Methyl RNA residue added** (2). Moreover, from a synthesis standpoint, the coupling efficiency of 2'-O-Methyl phosphoramidites are higher than those of RNA monomers, resulting in higher yield of full-length oligos.

Modifications Increasing Duplex Stability and Nuclease Resistance

Modification

Duplex Stability [ $T_m$  Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased

[genelink.com/oligo\\_modifications\\_reference/OMR\\_mod\\_category\\_intro.asp?mod\\_sp\\_cat\\_id=19](http://genelink.com/oligo_modifications_reference/OMR_mod_category_intro.asp?mod_sp_cat_id=19) >Click here for complete list of duplex stability modifications

### **ASO's and siRNA Modifications.**

Click this link to view ASO's and siRNA Modifications.

**ASO's and siRNA Delivery.** The development of effective delivery systems for antisense oligonucleotides is essential for their clinical therapeutic application. The most common delivery system involves a relatively hydrophobic molecule that can cross the lipid membrane. Cholesterol TEG, alpha-Tocopherol TEG ( a natural isomer of vitamin E), stearyl and GalNAc modifications have been shown to effective for delivery of ASO's and siRNA in addition to cell penetrating peptides. Click this link to view these modifications.

### **References**

1. Cotton, M.; Oberhauser, B.; Burnar, H. *et al.* 2'O methyl and 2'O ethyl oligoribonucleotides as inhibitors of the in vitro U7 snRNP-dependent messenger-RNA processing event. *Nucleic Acids Res.* (1991) , **19**:2629-2635.
2. Kawasaki, A.M. *et al.*, Uniformly modified 2'-deoxy-2'-fluoro phosphorothioate oligonucleotides as nuclease resistant antisense compounds with high affinity and specificity for RNA targets, *Journal of Medicinal Chemistry* (1993), **36**: 831-841.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 2'-O methyl U

Category	Nuclease Resistance
Modification Code	mU
Reference Catalog Number	27-6410U
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	320.2



Antisense Oligos (ODN) & siRNA Oligo Modifications

**Click here for more information on antisense modifications, design & applications.**

2'-O-Methyl bases are classified as a 2'-O-Methyl RNA monomer. 2'-O-Methyl nucleotides are most commonly used **to confer nuclease resistance** to an oligo designed for anti-sense, siRNA or aptamer-based research, diagnostic or therapeutic purposes, when specific 2'-OH is not required. Nuclease resistance can be further enhanced by phosphorothiolation of appropriate internucleotide linkages within the oligo.

The hydrogen bonding behavior of a 2'-O-Methyl RNA/RNA base pair is closer to that of an RNA/RNA base pair than a DNA/RNA base pair. Consequently, the presence of 2'-O-Methyl nucleotides **improves duplex stability**. Indeed, incorporation of a 2'-O-Methyl nucleotide into an anti-sense oligo (resulting in a 2'-O-Methyl RNA/DNA chimeric), lead to a **increase** in the T<sub>m</sub> of its duplex with RNA, relative to that formed by an unmodified anti-sense DNA oligo, **of 1.3°C per 2'-O-Methyl RNA residue added** (2). Moreover, from a synthesis standpoint, the coupling efficiency of 2'-O-Methyl phosphoramidites are higher than those of RNA monomers, resulting in higher yield of full-length oligos.

Modifications Increasing Duplex Stability and Nuclease Resistance

Modification

Duplex Stability [T<sub>m</sub> Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased

[genelink.com/oligo\\_modifications\\_reference/OMR\\_mod\\_category\\_intro.asp?mod\\_sp\\_cat\\_id=19](http://genelink.com/oligo_modifications_reference/OMR_mod_category_intro.asp?mod_sp_cat_id=19) >Click here for complete list of duplex stability modifications

### **ASO's and siRNA Modifications.**

Click this link to view ASO's and siRNA Modifications.

**ASO's and siRNA Delivery.** The development of effective delivery systems for antisense oligonucleotides is essential for their clinical therapeutic application. The most common delivery system involves a relatively hydrophobic molecule that can cross the lipid membrane. Cholesterol TEG, alpha-Tocopherol TEG ( a natural isomer of vitamin E), stearyl and GalNAc modifications have been shown to effective for delivery of ASO's and siRNA in addition to cell penetrating peptides. Click this link to view these modifications.

### **References**

1. Cotton, M.; Oberhauser, B.; Burnar, H. *et al.* 2'O methyl and 2'O ethyl oligoribonucleotides as inhibitors of the in vitro U7 snRNP-dependent messenger-RNA processing event. *Nucleic Acids Res.* (1991) , **19**:2629-2635.
2. Kawasaki, A.M. *et al.*, Uniformly modified 2'-deoxy-2'-fluoro phosphorothioate oligonucleotides as nuclease resistant antisense compounds with high affinity and specificity for RNA targets, *Journal of Medicinal Chemistry* (1993), **36**: 831-841.





## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 2-Amino dA

Category	Duplex Stability
Modification Code	2Am-dA
Reference Catalog Number	26-6525
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	328.22



2-Amino dA, Catalog#: 26-6525 has been discontinued effective June 25, 2025 and replaced by a more stable identical final modification of 2-6-diaminopurine (2-Amino dA) Catalog#: 26-6504. The original 2-Amino dA catalog number 26-6525 precursor base was labile and required mild synthesis conditions with low yield. The substitute is more stable.

#### 2-6-diaminopurine (2-Amino dA) Catalog: 26-6504-XX.

2-Amino-deoxyadenosine dA (also known as 2-6 diamino purine) can be used to **improve the ability of an oligo to hybridize to its target**. The 2-Amino-A nucleotide base forms three hydrogen bonds with thymine (T), compared with only two H-bonds between unmodified A and T. 2-Amino A:T base pairs thus have the same number of H-bonds as G:C base pairs do. Consequently, when a 2-Amino-dA oligo binds to its unmodified target, the T<sub>m</sub> of the duplex **is raised by 3oC per 2-Amino-dA residue added**, compared with the unmodified case (1). In addition, 2-Amino-A also destabilizes A-G wobble mismatches, presumably due to a steric clash between the 2-amino on A and the 2-amino on G. Thus 2-Amino-dA modified oligos show better specificity for a target than their unmodified counterparts.

Modifications Increasing Duplex Stability and Nuclease Resistance

Modification

Duplex Stability [T<sub>m</sub> Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased

[genelink.com/oligo\\_modifications\\_reference/OMR\\_mod\\_category\\_intro.asp?mod\\_sp\\_cat\\_id=19](http://genelink.com/oligo_modifications_reference/OMR_mod_category_intro.asp?mod_sp_cat_id=19) >Click here for complete list of duplex stability modifications

2-Amino dA has been shown to be particularly useful in the following oligo-based applications:

(a)Strong-binding PCR primers: 2-Amino-dA-modified PCR primers have been shown to prime far better than their unmodified counterparts in PCR reactions, consistently yielding more product per cycle, permitting amplification at very high annealing temperatures (as high as 72°C), and interestingly, allowing excellent priming from within palindromic sequences (1). The improvement in priming efficiency could significantly reduce the number of amplification-related mutations in PCR products. 2-Amino-dA primers also could be useful in several PCR applications, e.g., when short, specific primers are required, when only a limited quantity of template is available (e.g. ancient DNA), when DNA secondary structure in the primer binding site prevents binding of an unmodified primer, or when primer extension is blocked by downstream DNA secondary structure in the template.

(b)Selective Binding Complementary (SBC) Oligos: SBC oligos are complementary pairs of oligos that contain one or more modified base pairs (that is, each member of the pair is modified). Each individual modified base does not form a stable base pair with its modified partner, but does form a particularly stable base pair with its natural (unmodified) counterpart. Thus, two complementary SBC oligos do not form a stable duplex with each other, but each individual SBC oligo does form a very stable duplex with an unmodified complementary target. This property enables an SBC duplex to effectively bind with both the sense and anti-sense strands of a DNA or RNA duplex target. SBC oligos are useful as probes for investigation of branching secondary structures, and as anti-sense reagents against mRNA that has significant secondary structure.

An excellent pair of SBC oligos can be made by substituting 2-Amino-dA for A, and 2-Thio-dT for T. Because 2-Amino-dA only forms one hydrogen bond with 2-Thio-dT, these modified base pairs are very weak, and the corresponding duplex is unstable. However, both 2-Amino-dA and 2-Thio-dT bind effectively with T and A bases, respectively. In a classic study, SBC 20mers annealed against a DNA 20mer target exhibited T<sub>m</sub> values 10° higher than the corresponding DNA-DNA hybrid, whereas the SBC-SBC hybrid yielded T<sub>m</sub> values 30° lower (2). **References**

Lebedev, Y., *et al.* Oligonucleotides containing 2-aminoadenine and 5-methylcytosine are more effective as primers for PCR amplification than their unmodified counterparts. *Genetic Analysis: Biomolecular Engineering* (1996), **13**: 15-21.

Kutyavin, I.V.; Rhinehart, R.L.; Lukhtanov, E.A.; Gorn, V.V.; Meyer, R.B.; Gamper, H.B. *Oligonucleotides Containing 2-Aminoadenine and 2-Thiothymine Act as Selectively Binding Complementary Agents*. *Biochemistry* (1996), **35**: 11170-11176.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### 5-Me dC

Category	Epigenetics
Modification Code	5mdC
Reference Catalog Number	26-6413
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	303.21



5-methyl deoxycytosine (5-Me-dC) pairs with dG, and when substituted for dC in an oligonucleotide, increases the stability of the resulting duplex relative to the comparable unmodified form, raising the  $T_m$  by 1.3degC per 5-Me-dC residue added (1,2). 5-Me-dC thus can be used to **improve the ability of an oligo to hybridize to its target**. The presence of the hydrophobic 5-methyl group presumably acts to exclude water molecules from the duplex.  
Modifications Increasing Duplex Stability and Nuclease Resistance

#### Modification

#### Duplex Stability [ $T_m$ Increase]

Nuclease Resistance Locked Analog Bases Increased [2- 4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased Click here for complete list of duplex stability modifications

5-Me-dC is particularly useful in the following applications:

(a)Strong-binding PCR primers: 5-Me-dC-modified PCR primers have been shown to prime far better than their unmodified counterparts in PCR reactions, consistently yielding more product per cycle, permitting amplification at very high annealing temperatures (as high as 72degC), and interestingly, allowing excellent priming from within palindromic sequences (1). The improvement in priming efficiency could significantly reduce the number of amplification-related mutations in PCR products.

5-Me-dC primers also could be useful in several PCR applications, *e.g.*, when short, specific primers are required, when only a limited quantity of template is available (*e.g.* ancient DNA), when DNA secondary structure in the primer binding site prevents binding of an unmodified primer, or when primer extension is blocked by downstream DNA secondary structure in the template.

(b) **Anti-sense:** Anti-sense oligonucleotides containing a CpG motif induce pro-inflammatory responses after *in vivo* administration to animals, including human, via activation of Toll-like receptor 9 (TLR9). Substitution of 5-Me-dC for dC in these motifs can prevent or sharply reduce these undesirable immune responses (3).

(b) **DNA methylation studies:** Methylation of dC to 5-methyl-dC, when it occurs in CpG sites near promoters is associated with gene silencing, and is an important epigenetic mechanism in living organisms. Oligonucleotides incorporating 5-Me-dC have been used by a number of research groups as research tools to study the epigenetic effects of DNA methylation in such areas as tumorigenesis and the effects of cocaine on fetal heart development (4-6). **References**

1. Lebedev, Y.; Akopyants, N.; Azhikina, T.; Shevchenko, Y.; Potapov, V.; Stecenko, D.; Berg, D.; Sverdlov, E.. Oligonucleotides containing 2-aminoadenine and 5-methylcytosine are more effective as primers for PCR amplification than their nonmodified counterparts. *Genet Anal.* (1996), **13**: 15-21.
2. Xodo, L.E.; Manzini, G.; Quadrifoglio, F.; van der Marel, G.A.; van Boom, J.H. Effect of 5-methylcytosine on the stability of triple-stranded DNA—a thermodynamic study *Nucleic Acids Res.* (1991), **19**: 5625-5631.
3. Henry, S.P.; Stecker, K.; Brooks, D.; Monteith, D.; Conklin, B.; Bennett, C.F. Chemically modified oligonucleotides exhibit decreased immune stimulation in mice. *J. Pharmacol. Exp. Ther.* (2000), **292**: 468-479.
4. Mancini, D.N.; Singh, S.M.; Archer, T.K.; Rodenhiser, D.I. Site-specific DNA methylation in the neurofibromatosis (NF1) promoter interferes with binding of CREB and SP1 transcription factors. *Oncogene.* (1999), **18**: 4108-4119.
5. Zhang, H.; Darwanto, A.; Linkhart, T.A.; Sowers, L.C.; Zhang, L. Maternal Cocaine Administration Causes an Epigenetic Modification of Protein Kinase Ce Gene Expression in Fetal Rat Heart, *Mol. Pharmacol.* (2007), **71**: 1319-1328.
6. Ishii, T.; Fujishiro, M.; Masuda, M.; Teramoto, S.; Matsuse, T. A methylated oligonucleotide induced methylation of GSTP1 promoter and suppressed its expression in A549 lung adenocarcinoma cells. *Cancer Letters* (2004), **212**: 211-223.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Amino C6 U

Category	Minor Bases
Modification Code	AmC6U
Reference Catalog Number	27-6422
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	474.4





## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Amino dA C6

Category	Minor Bases
Modification Code	Am-dA-C6
Reference Catalog Number	26-6666
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	427.4





## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Amino dC C6

Category	Minor Bases
Modification Code	Am-dC-C6
Reference Catalog Number	26-6670
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	507.51







## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Amino dG C6

Category	Minor Bases
Modification Code	Am-dG-C6
Reference Catalog Number	26-6669
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	428.38







## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Amino dT C6

Category	Minor Bases
Modification Code	Am-dT-C6
Reference Catalog Number	26-6438
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	458.41





## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Cy3-5'

Category	Fluorescent Dyes
Modification Code	Cy3-5
Reference Catalog Number	26-6437
5 Prime	Y
3 Prime	N
Internal	N
Molecular Weight(mw)	630.78



[Click here for a list of fluorophores.](#)

Prices listed above are for 5' modification. Internal and 3' incurs additional charges and are post synthesis conjugation to a primary amino group thus an additional modification with an amino group is required. A C6 or C12 amino group can be placed at the 5' or for the 3' end a C3 or C7 amino and for internal positions an amino modified base is used, e.g Amino dT C6.

Cyanine 3 (Cy3) is a fluorescent dye that belongs to the Cyanine family of synthetic polymethine dyes. Cy3 is reactive, water-soluble, and has an absorbance maximum of 550 nm and an emission maximum of 570 nm. It is available as both a phosphoramidite and an NHS ester, and is used to fluorescently label oligonucleotides at either the 5' or 3' end, or internally. Cy3 plays a particularly important role in real-time PCR applications, being used as the reporter moiety in TaqMan probes (1), Scorpion primers (2) and Molecular Beacons (3). For such probes, Cy3 is most commonly paired with the dark quencher BHQ-2, as the two have excellent spectral overlap.

Cy3 can also be used to label DNA oligos for use as hybridization probes in other applications, such as Fluorescent In-Situ Hybridization (FISH). In 2010, Stoeckler and co-workers (4) reported that Cy3 double-labeling of FISH probes (at both ends) that were specific to ribosomal RNA targets in microorganisms at least doubles FISH signal intensity without affecting specificity. This Double Labeling of Oligonucleotide Probes for Fluorescence In Situ Hybridization (DOPE-FISH 0 strategy may provide an effective solution to the problem of low signal intensity, which is commonly observed when using corresponding singly-labeled FISH probes for microbe identification. As an added benefit, Cy3-doubly labeled probes were shown to increase the in situ accessibility of rRNA targets sites in microbes, which allows for greater probe design flexibility.

Applied Biosystems Proprietary Dyes & Possible Substitutions

Fluorophore

Color

Absorbance max (nm)

Emission max (nm) VIC Pink Red 538 554 Cal Orange 560 Pink Red 537 558 HEX Pink Red 535 556 NED Red Orange 546 575 Cy3 Red Orange 550 570 PET Red Orange 558 595 Cy3.

5 Red 588 604 ROX Red 575 602 Texas Red Red 583 603

**Click here for a list of fluorophores.**

#### **References**

1. Livak, K.J., Flood, S.J.A., Marmaro, J., Giusti, W., Deetz, K. Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. *PCR Methods Appl.* (1995), **4**: 1-6.
2. Thelwell, N., Millington, S., Solinas, A., Booth, J., Brown, T. Mode of action and application of Scorpion primers to mutation detection. *Nucleic Acids Res.* (2000), **28**: 3752-3761.
3. Tyagi, S., Kramer, F.R. Molecular beacons: probes that fluoresce upon hybridization. *Nat. Biotechnol.* (1996), **14**: 303-308.
4. Stoecker, K., Dorninger, C., Daims, H., Wagner, M. Double Labeling of Oligonucleotide Probes for Fluorescence In Situ Hybridization (DOPE-FISH) Improves Signal Intensity and Increases rRNA Accessibility. *Appl. Environ. Microb.* (2010), **76**: 922-926.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Cy3.5-5'

Category	Fluorescent Dyes
Modification Code	Cy3.5-5
Reference Catalog Number	26-6461
5 Prime	Y
3 Prime	N
Internal	N
Molecular Weight(mw)	607.7



Cyanine 3.5 (Cy3.5) is a fluorescent dye that belongs to the Cyanine family of synthetic polymethine dyes. Cy3.5 is reactive, water-soluble, and has an absorbance maximum of 581 nm and an emission maximum of 596 nm. It is available as a phosphoramidite, and is used to fluorescently label oligonucleotides at either the 5' or 3' end, or internally. Cy3.5 plays a particularly important role in real-time PCR applications, being used as a reporter moiety in TaqMan probes (1), Scorpion primers (2) and Molecular Beacons (3). For such probes, Cy3.5 is most commonly paired with the dark quencher BHQ-2, as the two have excellent spectral overlap.

Cy3 can also be used to label DNA oligos for use as hybridization probes in other applications, such as Fluorescent In-Situ Hybridization (FISH).

Applied Biosystems Proprietary Dyes & Possible Substitutions

Dye

Color

Absorbance max (nm)

Emission max (nm) VIC Pink Red 538 554 Cal Orange 560 Pink Red 537 558 HEX Pink Red 535 556 NED Red Orange 546 575 Cy3 Red Orange 550 570 PET Red Orange 558 595 Cy3.

5 Red 588 604 ROX Red 575 602 CAL Fluor Red 590 Red 569 591 Texas Red Red 583 603

**Click here for a list of fluorophores.**

#### **References**

1. Livak, K.J., Flood, S.J.A., Marmaro, J., Giusti, W., Deetz, K. Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. *PCR Methods Appl.* (1995), **4**: 1-6.
2. Thelwell, N., Millington, S., Solinas, A., Booth, J., Brown, T. Mode of action and application of Scorpion primers to mutation detection. *Nucleic Acids Res.* (2000), **28**: 3752-3761.
3. Tyagi, S., Kramer, F.R. Molecular beacons: probes that fluoresce upon hybridization. *Nat. Biotechnol.* (1996), **14**: 303-308.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Cy5-5'

Category	Fluorescent Dyes
Modification Code	Cy5-5
Reference Catalog Number	26-6436
5 Prime	Y
3 Prime	N
Internal	N
Molecular Weight(mw)	656.81



Cy5-5' [26-6436-XX]

[Click here for a list of fluorophores.](#)

Prices listed above are for 5' modification. Internal and 3' incurs additional charges and are post synthesis conjugation to a primary amino group thus an additional modification with an amino group is required. A C6 or C12 amino group can be placed at the 5' or for the 3' end a C3 or C7 amino and for internal positions an amino modified base is used, e.g Amino dT C6.

Cyanine 5 (Cy5) is a fluorescent dye that belongs to the Cyanine family of synthetic polymethine dyes. Cy5 is reactive, water-soluble, and has an absorbance maximum of 649 nm and an emission maximum of 670 nm. It is available as both a phosphoramidite and an NHS ester, and is used to fluorescently label oligonucleotides at either the 5' or 3' end, or internally. Cy5 plays a particularly important role in real-time PCR applications, being used as a reporter moiety in TaqMan probes (1), Scorpion primers (2) and Molecular Beacons (3). For such probes, Cy5 is most commonly paired with the dark quencher BHQ-3, as the two have excellent spectral overlap.

Cy5 can also be used to label DNA oligos for use as hybridization probes in other applications, such as Fluorescent In-Situ Hybridization (FISH). In 2010, Stoeckler and co-workers (4) reported that Cy5 double-labeling of FISH probes (at both ends) that were specific to ribosomal RNA targets in microorganisms at least doubles FISH signal intensity without affecting specificity. This Double Labeling of Oligonucleotide Probes for Fluorescence In Situ Hybridization (DOPE-FISH) strategy may provide an effective solution to the problem of low signal intensity, which is commonly observed when using corresponding singly-labeled FISH probes for microbe identification. As an added benefit, Cy5-doubly labeled probes were shown to increase the in situ accessibility of rRNA targets sites in microbes, which allows for greater probe design flexibility.

[Click here for list of quenchers.](#)

[Click here for a list of fluorophores.](#)

## Near Infrared Fluorophore Spectral Data & Quencher Selection Guide

Fluorophore Name

Absorbance Max, nm +/-10

Emission Max, nm +/-10

Extinction Coefficient\*

Color\*\*

Quencher

**Cy5 650 665 250,000**

**IRDye 650 NHS 650 665 230,000**

**AZ647 NHS 655 680 191,800**

**AZ680 NHS 678 701 185,000**

**Cy5.5 684 710 198,000**

**IRDye 700 NHS 684 710 288,000**

**AZdye700 NHS 696 719 192,000**

**Atto 700 NHS 700 716 120,000**

**Atto 725 NHS 728 751 120,000**

**Atto 740 NHS 743 763 120,000**

**Cy7 NHS 740 773 199,000**

**IRDye 750 NHS 756 776 260,000**

**cy7.5 NHS 788 808 223,000**



IRDye 800 NHS 795 819 240,000

\* Extinction coefficient at  $\lambda$  (max) in cm<sup>-1</sup>M<sup>-1</sup>. \*\* Typical emission color seen through the eyepiece of a conventional fluorescence microscope with appropriate filters. Near-IR region. Human vision is insensitive to light beyond ~650 nm; it is not possible to view near-IR fluorescent dyes.

[Click here for a list of fluorophores.](#)

[Click here for list of quenchers.](#)

#### References

1. Livak, K.J., Flood, S.J.A., Marmaro, J., Giusti, W., Deetz, K. Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. *PCR Methods Appl.* (1995), 4: 1-6.
2. Thelwell, N., Millington, S., Solinas, A., Booth, J., Brown, T. Mode of action and application of Scorpion primers to mutation detection. *Nucleic Acids Res.* (2000), 28: 3752-3761.
3. Tyagi, S., Kramer, F.R. Molecular beacons: probes that fluoresce upon hybridization. *Nat. Biotechnol.* (1996), 14: 303-308.
4. Stoecker, K., Dorninger, C., Daims, H., Wagner, M. Double Labeling of Oligonucleotide Probes for Fluorescence In Situ Hybridization (DOPE-FISH) Improves Signal Intensity and Increases rRNA Accessibility. *Appl. Environ. Microb.* (2010), 76: 922-926.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Cy5.5

Category	Fluorescent Dyes
Modification Code	Cy5.5-5
Reference Catalog Number	26-6460
5 Prime	Y
3 Prime	N
Internal	N
Molecular Weight(mw)	633.74



Cyanine 5.5 (Cy5.5) is a far-red/near infra red (NIR) fluorescent dye that belongs to the Cyanine family of synthetic polymethine dyes. Cy5.5 is reactive, water-soluble, and has an absorbance maximum of 675 nm and an emission maximum of 694 nm. It is available as a phosphoramidite, and is used to fluorescently label oligonucleotides at either the 5' or 3' end, or internally. Cy5.5 plays a particularly important role in real-time PCR applications, being used as a reporter moiety in TaqMan probes (1), Scorpion primers (2) and Molecular Beacons (3). For such probes, Cy5.5 is most commonly paired with the dark quencher BHQ-3, as the two have excellent spectral overlap.

Cy5.5 can also be used to label DNA oligos for use as hybridization probes in other applications, such as Fluorescent In-Situ Hybridization (FISH).

#### Near Infrared Fluorophore Spectral Data & Quencher Selection Guide

Fluorophore Name

Absorbance Max, nm +/-10

Emission Max, nm +/-10

Extinction Coefficient\*

Color\*\*

Quencher

**Cy5 650 665 250,000**

**IRDye 650 NHS 650 665 230,000**

asp?modid=516">AZ647 NHS 655 680 191,800

**AZ680 NHS 678 701 185,000**

**Cy5.5 684 710 198,000**

**IRDye 700 NHS 684 710 288,000**

**AZdye700 NHS 696 719 192,000**

**Atto 700 NHS 700 716 120,000**

**Atto 725 NHS 728 751 120,000**

**Atto 740 NHS 743 763 120,000**

Cy7 NHS 740 773 199,000

IRDye 750 NHS 756 776 260,000

cy7.5 NHS 788 808 223,000

IRDye 800 NHS 795 819 240,000

\* Extinction coefficient at  $\lambda$  (max) in cm<sup>-1</sup>M<sup>-1</sup>. \*\* Typical emission color seen through the eyepiece of a conventional fluorescence microscope with appropriate filters. Near-IR region. Human vision is insensitive to light beyond ~650 nm; it is not possible to view near-IR fluorescent dyes.

[Click here for a list of fluorophores.](#)

[Click here for list of quenchers.](#)

#### References

1. Livak, K.J., Flood, S.J.A., Marmaro, J., Giusti, W., Deetz, K. Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. *PCR Methods Appl.* (1995), 4: 1-6.
2. Thelwell, N., Millington, S., Solinas, A., Booth, J., Brown, T. Mode of action and application of Scorpion primers to mutation detection. *Nucleic Acids Res.* (2000), 28: 3752-3761.
3. Tyagi, S., Kramer, F.R. Molecular beacons: probes that fluoresce upon hybridization. *Nat. Biotechnol.* (1996), 14: 303-308.
4. Stoecker, K., Dorninger, C., Daims, H., Wagner, M. Double Labeling of Oligonucleotide Probes for Fluorescence In Situ Hybridization (DOPE-FISH) Improves Signal Intensity and Increases rRNA Accessibility. *Appl. Environ. Microb.* (2010), 76: 922-926.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Cy7 NHS

Category	Fluorescent Dyes
Modification Code	Cy7-N
Reference Catalog Number	26-6474
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	682



**Click here for a list of fluorophores.**

This modification is a post synthesis conjugation to a primary amino group thus an additional modification with an amino group is required. A C3, C6 or C12 amino group can be placed at the 5' or for the 3' end a C3 or C7 amino and for internal positions an amino modified base is used, e.g Amino dT C6.

Yield of Post Synthesis NHS, Maleimide & Click Ligand Conjugation\* Oligo Scale of Synthesis Yield, nmols 50 nmol 2 nmol 200 nmol 5 nmol 1 umol 16 nmol 2 umol 30 nmol 5 umol 75 nmol 10 umol 150 nmol 15 umol 225 nmol \* The yield will be lower for oligos longer than 50mer. Click here for yield table of long oligos. \* Click here for RNA Oligos scale of synthesis and yield. **NHS Ligand conjugation** requires a primary amino group. Gene Link offers a wide selection of amino modifications for 5', 3' and internal sites. Click here for a list of conjugation chemistry modifications. **Maleimide Ligand conjugation** requires a thiol group. Gene Link offers a wide selection of thiol modifications for 5', 3' and internal sites.

Click here for a list of conjugation chemistry modifications. **Click Chemistry Ligand conjugation** requires a corresponding Click modification; examples Alkyne:Azide, Azide:DBCO, BCN:Azide, BCN: TCO:Tetrazine. Gene Link offers a wide selection of click modifications for 5', 3' and internal sites. Click here for a list of click chemistry modifications.

#### Near Infrared Fluorophore Spectral Data & Quencher Selection Guide

Fluorophore Name

Absorbance Max, nm +/-10

Emission Max, nm +/-10

Extinction Coefficient\*

Color\*\*

Quencher

**Cy5 650 665 250,000**

**IRDye 650 NHS 650 665 230,000**

**AZ647 NHS 655 680 191,800**

**AZ680 NHS 678 701 185,000**

**Cy5.5 684 710 198,000**

**IRDye 700 NHS 684 710 288,000**

**AZdye700 NHS 696 719 192,000**

**Atto 700 NHS 700 716 120,000**

**Atto 725 NHS 728 751 120,000**

**Atto 740 NHS 743 763 120,000**

**Cy7 NHS 740 773 199,000**

**IRDye 750 NHS 756 776 260,000**

**cy7.5 NHS 788 808 223,000**

IRDye 800 NHS 795 819 240,000

\* Extinction coefficient at  $\lambda$  (max) in  $\text{cm}^{-1}\text{M}^{-1}$ . \*\* Typical emission color seen through the eyepiece of a conventional fluorescence microscope with appropriate filters. Near-IR region. Human vision is insensitive to light beyond ~650 nm; it is not possible to view near-IR fluorescent dyes.

[Click here for a list of fluorophores.](#)

[Click here for list of quenchers.](#)

Cyanine 7(Cy7) NHS ester is a fluorescent dye that belongs to the Cyanine family of synthetic polymethine dyes. Cy7 is reactive, water-soluble, and has an absorbance maximum of 747 nm and an emission maximum of 776 nm, which is in the near IR. It is available as an NHS ester, and is used to fluorescently label oligonucleotides at either the 5' or 3' end, or internally. Because it is a near IR dye, Cy7 has very little background fluorescence associated with it (1). It is thus an excellent choice for labeling oligo probes slated for in vivo applications, because the minimal scattering and absorption of near-IR photons by cellular tissue ensures higher S/N ratio, and better sensitivity. For example, Fluorescent Resonance Energy Transfer (FRET) oligonucleotide duplexes using Cy5.5 as the donor on one strand and Cy7 as the acceptor on the complementary strand have been used to detect and characterize transcription factor NF-kappaB p50 protein binding to DNA (2)

**Caution:** Cy7 is intensely colored and very reactive. Care should be exercised when handling the vial containing the C7-labeled oligo to avoid staining clothing, skin, and other items. Also, because Cy7 is in the form of an NHS ester, the oligo first must be synthesized with an Amino C6 Linker (for the ends) or the Amino C6 version of the base phosphoramidite (for internal labeling). The Cy7-NHS ester is then manually attached to the oligo through the amino group in a separate reaction post-synthesis.

#### References

1. Benson, R.C., Kues, H.A. Absorption and Fluorescence Properties of Cyanine Dyes. *J. Chem. Eng. Data* (1977), 22: 379-383.
2. Zhang, S., Metelev, V., Tabatadze, D., Zamecnik, P.C., Bogdanov, A. Fluorescence resonance energy transfer in near-infrared fluorescent oligonucleotide probes for detecting protein-DNA interactions. *Proc. Nat. Acad. Sci. USA*. (2008), 105: 4156-4161.





## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Fam dT

Category	Fluorescent Dyes
Modification Code	Fam-dT
Reference Catalog Number	26-6422
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	815.71





## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Fluorescein-5'

Category	Fluorescent Dyes
Modification Code	FI-5
Reference Catalog Number	26-6426
5 Prime	Y
3 Prime	N
Internal	N
Molecular Weight(mw)	567.6



Fluorescein-5' FITC (Fluorescein isothiocyanate) 5'  
[26-6426-XX]



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Hex-5'

Category	Fluorescent Dyes
Modification Code	Hex-5
Reference Catalog Number	26-6432
5 Prime	Y
3 Prime	N
Internal	N
Molecular Weight(mw)	744.13



**Click here for a list of fluorophores.**

Hexachloro-fluorescein (HEX) is hexachlorinated version of the fluorescent dye fluorescein, and is used for labeling oligonucleotides at either the 5' or 3' end. HEX has an absorbance maximum of 535 nm and an emission maximum of 556 nm. HEX can be used in real-time PCR applications as a reporter moiety in TaqMan probes (1), Scorpion primers (2) and Molecular Beacons (3). For such probes, HEX is most commonly paired with the dark quencher BHQ-1, as the two have good spectral overlap.

HEX also can be used to label DNA oligos for use as hybridization probes in a variety of in vivo and in vitro research or diagnostic applications, as well as for structure-function studies of DNA, RNA, and protein-oligonucleotide complexes. Oligos labeled with HEX at the 5' end can be used as PCR and DNA sequencing primers to generate fluorescently-labeled PCR, sequencing or genetic analysis (AFLP or microsatellite) products. NOTE: If HEX is on the 3' end of the oligo, it cannot be used as a primer in PCR-based applications.

Applied Biosystems Proprietary Dyes & Possible Substitutions

Dye

Color

Absorbance max (nm)

Emission max (nm) VIC Pink Red 538 554 Cal Orange 560 Pink Red 537 558 HEX Pink Red 535 556 NED Red Orange 546 575 Cy3 Red Orange 550 570 PET Red Orange 558 595 Cy3.

5 Red 588 604 ROX Red 575 602 Texas Red Red 583 603

**Click here for a list of fluorophores.**

#### **References**

1. Livak, K.J., Flood, S.J.A., Marmaro, J., Giusti, W., Deetz, K. Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. *PCR Methods Appl.* (1995), **4**: 1-6.
2. Thelwell, N., Millington, S., Solinas, A., Booth, J., Brown, T. Mode of action and application of Scorpion primers to mutation detection. *Nucleic Acids Res.* (2000), **28**: 3752-3761.
3. Tyagi, S., Kramer, F.R. Molecular beacons: probes that fluoresce upon hybridization. *Nat. Biotechnol.* (1996), **14**: 303-308.



## Product Specifications


Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

## Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### Tet-5'

Category	Fluorescent Dyes
Modification Code	Tet-5
Reference Catalog Number	26-6433
5 Prime	Y
3 Prime	N
Internal	N
Molecular Weight(mw)	675.24



**5'-Tetrachloro-Fluorescein (TET)**  
**[26-6433-XX]**