



Product Specifications

Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

AZDye-680-N

Category	Fluorescent Dyes
Modification Code	AZDye-680-N
Reference Catalog Number	26-6716
5 Prime	Y
3 Prime	Y
Internal	Y
Molecular Weight(mw)	924.21

Click here for a list of fluorophores.

This modification is a post synthesis conjugation to a primary amino group thus an additional modification with an amino group is required. A C3, C6 or C12 amino group can be placed at the 5' or for the 3' end a C3 or C7 amino and for internal positions an amino modified base is used, e.g Amino dT C6.

Yield of Post Synthesis NHS, Maleimide & Click Ligand Conjugation* Oligo Scale of Synthesis Yield, nmols 50 nmol 2 nmol 200 nmol 5 nmol 1 umol 16 nmol 2 umol 30 nmol 5 umol 75 nmol 10 umol 150 nmol 15 umol 225 nmol * The yield will be lower for oligos longer than 50mer. Click here for yield table of long oligos. * Click here for RNA Oligos scale of synthesis and yield. **NHS Ligand conjugation** requires a primary amino group. Gene Link offers a wide selection of amino modifications for 5', 3' and internal sites. Click here for a list of conjugation chemistry modifications. **Maleimide Ligand conjugation** requires a thiol group. Gene Link offers a wide selection of thiol modifications for 5', 3' and internal sites.

Click here for a list of conjugation chemistry modifications. **Click Chemistry Ligand conjugation** requires a corresponding Click modification; examples Alkyne:Azide, Azide:DBCO, BCN:Azide, BCN: TCO:Tetrazine. Gene Link offers a wide selection of click modifications for 5', 3' and internal sites. Click here for a list of click chemistry modifications.

Human vision is insensitive to light beyond ~650 nm; it is not possible to view near-IR fluorescent dyes.

AZDyes are a set of fluorescent dyes that span the visible electromagnetic spectrum, as well as some of the near-IR. The absorbance range is 346-749 nm, and the emission range is 442-775 nm. Generally speaking, AZDyes are brighter, chemically more stable, and less pH-sensitive than other fluorescent dyes commonly used to label oligonucleotides (1). Because they currently only are in the form of NHS esters, oligos first must be synthesized with an Amino Linker modification (either at the ends or internally). The appropriate AZDye is then manually attached to the oligo through the amino group in a separate reaction post-synthesis. The list of currently available dyes includes AZDye 350, -405, -430, -488, -500, -514, -532, -546, -555, -568, -594, -610, -633, -647, -660, -680, -700, -750, with the number indicating the appropriate absorbance wavelength for the particular dye. AZDyes are suitable for a variety of in vitro and in vivo applications. However, for in vivo experiments, users should note that dyes 350/405, being "blue" dyes, require higher-energy excitations than the other. Users of these particular dyes should confirm that the higher-energy required for excitation does not damage the relevant cells or tissues being used in the in vivo experiments.

-

Near Infrared Fluorophore Spectral Data & Quencher Selection Guide

Fluorophore Name

Absorbance Max, nm +/-10

Emission Max, nm +/-10

Extinction Coefficient*

Color**

Quencher

Cy5 650 665 250,000

IRDye 650 NHS 650 665 230,000

AZ647 NHS 655 680 191,800

AZ680 NHS 678 701 185,000

Cy5.5 684 710 198,000

IRDye 700 NHS 684 710 288,000

AZdye700 NHS 696 719 192,000

Atto 700 NHS 700 716 120,000

Atto 725 NHS 728 751 120,000

Atto 740 NHS 743 763 120,000

Cy7 NHS 740 773 199,000

IRDye 750 NHS 756 776 260,000

cy7.5 NHS 788 808 223,000

IRDye 800 NHS 795 819 240,000

* Extinction coefficient at λ (max) in $\text{cm}^{-1}\text{M}^{-1}$. ** Typical emission color seen through the eyepiece of a conventional fluorescence microscope with appropriate filters. Near-IR region. Human vision is insensitive to light beyond ~650 nm; it is not possible to view near-IR fluorescent dyes.

[Click here for a list of fluorophores.](#)

[Click here for list of quenchers.](#)

References

1. Didenko, V.V. DNA Probes Using Fluorescence Resonance Energy Transfer (FRET): Designs and Applications. *Biotechniques* (2001), 31: 1106-1121.
2. Livak, K.J., Flood, S.J.A., Marmaro, J., Giusti, W., Deetz, K. Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. *PCR Methods Appl.* (1995), 4: 1-6.
3. Thelwell, N., Millington, S., Solinas, A., Booth, J., Brown, T. Mode of action and application of Scorpion primers to mutation detection. *Nucleic Acids Res.* (2000), 28: 3752-3761.
4. Tyagi, S., Kramer, F.R. Molecular beacons: probes that fluoresce upon hybridization. *Nat. Biotechnol.* (1996), 14: 303-308.