


## Product Specifications

Custom Oligo Synthesis, antisense oligos, RNA oligos, chimeric oligos, Fluorescent dyes, Affinity Ligands, Spacers & Linkers, Duplex Stabilizers, Minor bases, labeled oligos, Molecular Beacons, siRNA, phosphonates Locked Nucleic Acids (LNA); 2'-5' linked Oligos

# Oligo Modifications

For research use only. Not for use in diagnostic procedures for clinical purposes.

### propyne dU



It has been shown that C-5 propyne analogs of dC and dT when substituted in phosphorothioate oligonucleotide imparts greater inhibition of gene expression due to increased binding affinity to the target mRNA and increased stability. Based on the above information antisense oligonucleotide could either be Phosphorothioated at all diester linkages or combined with substitutions of dC and dT by C-5 propyne analogs pdC and pdU. Modifications Increasing Duplex Stability and Nuclease Resistance

### Modification

### Duplex Stability [Tm Increase]

Nuclease Resistance Locked Analog Bases Increased [2-4C per substitution] Increased 2-Amino-dA Increased [3.0C per substitution] Similar to DNA C-5 propynyl-C Increased [2.8C per substitution] Increased C-5 propynyl-U Increased [1.7C per substitution] Increased 2'-Fluoro Increased [1.8C per substitution] Increased 5-Methyl-dC Increased [1.3C per substitution] Similar to DNA 2'-O Methyl Increased Increased Phosphorothioate Slightly decreased Increased Click here for complete list of duplex stability modifications The use of propyne analogs is covered by patents and licensing agreements. The sale of propyne-modified oligos is for research use only. See license agreement\*

